

ORDER OF THE RUNGE–KUTTA METHOD AND EVOLUTION OF THE STABILITY REGION¹

Hippolyte Séka[†], Kouassi Richard Assui^{††}

Institut National Polytechnique Houphouët–Boigny,
BP 1093 Yamoussoukro, Côte d’Ivoire

[†]hippolyte.seka@inphb.ci, ^{††}r.assui@yahoo.fr

Abstract: In this article, we demonstrate through specific examples that the evolution of the size of the absolute stability regions of Runge–Kutta methods for ordinary differential equation does not depend on the order of methods.

Keywords: Stability region, Runge–Kutta methods, Ordinary differential equations, Order of methods.

Introduction

Representations of the stability regions of Runge–Kutta methods are presented in several literatures [1–8, 11, 13]. It has been found that the stability region varies according to the order of the method. However, it is not proven in the literature whether or not there is a relation between the evolution of the size of the region of stability and the order of the method. In this article, we demonstrate that the evolution of the size of the stability region does not depend on the order of the methods. For that we exhibit methods whose regions of stability grow according to the order. Subsequently, we give a counter-example where we introduce a new 8 order method [12]. We compare the stability region of this new 8 order method with those of certain lower order methods. We show that the stability regions of lower order methods are larger than that of the new 8 order method. The study will be done in accordance with the following plan: in Section 2 we describe some generalities on the stability regions, in Section 3 we present some stability functions, in Section 4 we present the new 8 order method and its stability regions, Section 5 we give a conclusion.

1. Generalities on the stability regions

Consider a general form of the first-order ODE given below:

$$y' = f(x, y(x)), \tag{1.1}$$

with the initial condition $y(x_0) = y_0$ for the interval $x_0 \leq x \leq x_n$. Here, x is the independent variable, y is the dependent variable, n is the number of point values, and f is the function of the derivation. The goal is to determine the unknown function $y(x)$ whose derivative satisfies (1.1) and the corresponding initial values. In doing so, let us discretize the interval $x_0 \leq x \leq x_n$ to be

$$x_0, \quad x_1 = x_0 + h, \quad x_2 = x_0 + 2h, \dots, \quad x_n = x_0 + nh,$$

¹ We would like to express our deepest appreciation and gratitude to Professor Sergey Khashin of Ivanovo State University who provided us the possibility to coordinate and complete this article.

where h is the fixed step size. With the initial condition $y(x_0) = y_0$, the unknown grid function $y_1, y_2, y_3, \dots, y_n$ can be calculated by using the Runge–Kutta method of the order 8 (RK8 method).

The 8-th order method is thus obtained by the resolution of the 200 equations with 11 stages [12] on Maple.

Lets consider the Butcher tableau of 8 order and 11 steps RK method (see Fig. 1):

0											
c_2	$a_{2,1}$										
c_3	$a_{3,1}$	$a_{3,2}$									
c_4	$a_{4,1}$	$a_{4,2}$	$a_{4,3}$								
c_5	$a_{5,1}$	$a_{5,2}$	$a_{5,3}$	$a_{5,4}$							
c_6	$a_{6,1}$	$a_{6,2}$	$a_{6,3}$	$a_{6,4}$	$a_{6,5}$						
c_7	$a_{7,1}$	$a_{7,2}$	$a_{7,3}$	$a_{7,4}$	$a_{7,5}$	$a_{7,6}$					
c_8	$a_{8,1}$	$a_{8,2}$	$a_{8,3}$	$a_{8,4}$	$a_{8,5}$	$a_{8,6}$	$a_{8,7}$				
c_9	$a_{9,1}$	$a_{9,2}$	$a_{9,3}$	$a_{9,4}$	$a_{9,5}$	$a_{9,6}$	$a_{9,7}$	$a_{9,8}$			
c_{10}	$a_{10,1}$	$a_{10,2}$	$a_{10,3}$	$a_{10,4}$	$a_{10,5}$	$a_{10,6}$	$a_{10,7}$	$a_{10,8}$	$a_{10,9}$		
c_{11}	$a_{11,1}$	$a_{11,2}$	$a_{11,3}$	$a_{11,4}$	$a_{11,5}$	$a_{11,6}$	$a_{11,7}$	$a_{11,8}$	$a_{11,9}$	$a_{11,10}$	
	b_1	b_2	b_3	b_4	b_5	b_6	b_7	b_8	b_9	b_{10}	b_{11}

Figure 1. Butcher tableau of RK8 method.

The numerical solution is given by the formula

$$y_{i+1} = y_i + h \left(\sum_{s=1}^{11} b_s k_s \right), \quad (1.2)$$

with

$$k_s = f \left(x_i + c_s h, y_i + h \sum_{j=1}^{s-1} a_{s,j} k_j \right), \quad x_{i+1} = x_i + h. \quad (1.3)$$

The concept of absolute stability, in its simplest form, is based on the analysis of the behavior, according to the values of the step h , of the numerical solutions of the model equation [9–12]:

$$u'(t) = \lambda u(t). \quad (1.4)$$

Using (1.3) and (1.4), we obtain:

$$\text{for } s \geq 1, \quad k_s = \lambda \left(y_i + h \sum_{j=1}^{s-1} a_{s,j} k_j \right);$$

which gives:

$$y_{i+1} = \zeta(h\lambda) y_i.$$

If $z = h\lambda$, then the absolute stability region is the set

$$\{z \in \mathbb{C} \mid |\zeta(z)| \leq 1\}.$$

2. Presentation of some stability functions

Consider the standard Runge-Kutta methods of orders 1 to 4. When (1.2) and (1.3) are applied to the model problem (1.4), the resulting equations are

$$\text{RK1:} \quad y_{i+1} = (1 + z) y_i;$$

$$\text{RK2:} \quad y_{i+1} = \left(1 + z + \frac{z^2}{2}\right) y_i;$$

$$\text{RK3:} \quad y_{i+1} = \left(1 + z + \frac{z^2}{2} + \frac{z^3}{6}\right) y_i;$$

$$\text{RK4:} \quad y_{i+1} = \left(1 + z + \frac{z^2}{2} + \frac{z^3}{6} + \frac{z^4}{24}\right) y_i.$$

The stability regions are shown at the next figure:




Figure 2. Evolution of the stability region according to the order.

We can see the evolution of the size of the region of stability as the order of the method increases. Let's now give a counterexample for which the stability region is very small.

3. Presentation of the new 8 order method and its stability regions

The family of the 8th order method is thus obtained by the resolution of the 200 equations with 11 stages [12] on Maple. This method depends on free parameters b_8 and $a_{10,5}$ [12].

Some of related coefficients have fixed values, not depending on b_8 and $a_{10,5}$, these coefficients are:

$$\begin{aligned}
b_1 &= \frac{1}{20}; & b_2 &= 0; & b_3 &= 0; & b_4 &= 0; & b_5 &= 0; & b_6 &= 0; & b_9 &= \frac{16}{45}; & b_{10} &= \frac{49}{180}; & b_{11} &= \frac{1}{20}; \\
c_2 &= \frac{1}{2}; & c_3 &= \frac{1}{2}; & c_4 &= \frac{7 + \sqrt{21}}{14}; & c_5 &= \frac{7 + \sqrt{21}}{14}; & c_6 &= \frac{1}{2}; \\
c_7 &= \frac{7 - \sqrt{21}}{14}; & c_8 &= \frac{7 - \sqrt{21}}{14}; & c_9 &= \frac{1}{2}; & c_{10} &= \frac{7 + \sqrt{21}}{14}; & c_{11} &= 1; \\
a_{2,1} &= \frac{1}{2}; \\
a_{3,1} &= \frac{1}{4}; & a_{3,2} &= \frac{1}{4}; \\
a_{4,1} &= \frac{1}{7}; & a_{4,2} &= \frac{-7 - 3\sqrt{21}}{98}; & a_{4,3} &= \frac{21 + 5\sqrt{21}}{49}; \\
a_{5,1} &= \frac{11 + \sqrt{21}}{84}; & a_{5,2} &= 0; & a_{5,3} &= \frac{4\sqrt{21}}{63} + \frac{2}{7}; & a_{5,4} &= \frac{21 - \sqrt{21}}{252}; \\
a_{6,1} &= \frac{5 + \sqrt{21}}{48}; & a_{6,2} &= 0; & a_{6,3} &= \frac{9 + \sqrt{21}}{36}; & a_{6,4} &= \frac{-231 + 14\sqrt{21}}{360}; & a_{6,5} &= \frac{63 - 7\sqrt{21}}{80}; \\
a_{7,1} &= \frac{10 - \sqrt{21}}{42}; & a_{7,2} &= 0; \\
a_{9,1} &= \frac{1}{32}; & a_{9,2} &= 0; \\
a_{10,1} &= \frac{1}{14}; & a_{10,2} &= 0; & a_{10,9} &= \frac{4\sqrt{21}}{35} + \frac{132}{245}; \\
a_{11,1} &= 0; & a_{11,2} &= 0; & a_{11,9} &= \frac{28 - 28\sqrt{21}}{45}; & a_{11,10} &= \frac{49 - 7\sqrt{21}}{18}.
\end{aligned}$$

And the others are expressed in terms of b_8 and $a_{10,5}$:

$$\begin{aligned}
b_7 &= -b_8 + \frac{49}{180}; \\
a_{7,3} &= -(24/35)a_{10,5} - 136/105 - (12/245)a_{10,5}\sqrt{21} + (656/2205)\sqrt{21}; \\
a_{7,4} &= 7 - (3/10)a_{10,5}\sqrt{21} - (71/45)\sqrt{21} + (3/10)a_{10,5}; \\
a_{7,5} &= -(3/10)a_{10,5} + (3/10)a_{10,5}\sqrt{21} - 43/6 + (169/105)\sqrt{21}; \\
a_{7,6} &= -(277/735)\sqrt{21} + 181/105 + (12/245)a_{10,5}\sqrt{21} + (24/35)a_{10,5}; \\
a_{8,1} &= -\frac{180b_8\sqrt{21} - 49\sqrt{21} - 1800b_8 + 343}{7560b_8}; & a_{8,2} &= 0; \\
a_{8,5} &= -\frac{441a_{10,5}\sqrt{21} - 3240a_{7,5}b_8 - 28\sqrt{21} + 882a_{7,5} - 2205a_{10,5} + 147}{3240b_8}; \\
a_{8,6} &= \frac{72a_{10,5}\sqrt{21} + 1620a_{7,6}b_8 - 29\sqrt{21} - 441a_{7,6} - 252a_{10,5} + 119}{1620b_8};
\end{aligned}$$

And also:

$$\begin{aligned}
a_{8,3} &= -\frac{900b_8\sqrt{21} + 11340a_{7,2}b_8 + 11340a_{8,6}b_8 - 98\sqrt{21} - 3087a_{7,2} - 4860b_8 + 686}{11340b_8}; \\
a_{8,7} &= \frac{49}{1620b_8}; \\
a_{8,4} &= \frac{(c_8^2/2) - a_{8,2}c_2 - a_{8,3}c_3 - a_{8,5}c_5 - a_{8,6}c_6 - a_{8,7}c_7}{c_4}; \\
a_{9,3} &= (1/8)a_{10,5}\sqrt{21} - (1/8)a_{10,5} - (1/72)\sqrt{21} + 1/72; \\
a_{9,4} &= -49/288 - (7/32)a_{10,5}\sqrt{21} + (7/288)\sqrt{21} + (49/32)a_{10,5}; \\
a_{9,5} &= (7/32)a_{10,5}\sqrt{21} - (35/576)\sqrt{21} - (49/32)a_{10,5} + 21/64; \\
a_{9,6} &= -(1/8)a_{10,5}\sqrt{21} + (1/8)a_{10,5} + (1/72)\sqrt{21} + 5/36; \\
a_{9,7} &= 91/576 + (7/192)\sqrt{21} - (585/1568)b_8\sqrt{21} - (405/224)b_8; \\
a_{9,8} &= (585/1568)b_8\sqrt{21} + (405/224)b_8; \\
a_{10,3} &= -(6/49)a_{10,5}\sqrt{21} - (2/7)a_{10,5} + (2/147)\sqrt{21} + 2/63; \\
a_{10,4} &= 1/9 - a_{10,5}; \\
a_{10,6} &= (2/7)a_{10,5} - 803/2205 + (6/49)a_{10,5}\sqrt{21} - (59/735)\sqrt{21}; \\
a_{10,7} &= 1/9 + (1/42)\sqrt{21} + (2295/686)b_8 + (495/686)b_8\sqrt{21}; \\
a_{10,8} &= -(2295/686)b_8 - (495/686)b_8\sqrt{21}; \\
a_{11,3} &= (2/3)a_{10,5}\sqrt{21} - (2/3)a_{10,5} - (2/27)\sqrt{21} + 2/27; \\
a_{11,4} &= -(7/6)a_{10,5}\sqrt{21} + (7/54)\sqrt{21} + (49/6)a_{10,5} - 49/54; \\
a_{11,5} &= (7/27)\sqrt{21} - 77/54 - (49/6)a_{10,5} + (7/6)a_{10,5}\sqrt{21}; \\
a_{11,6} &= (2/3)a_{10,5} - 64/135 - (2/3)a_{10,5}\sqrt{21} + (94/135)\sqrt{21}; \\
a_{11,7} &= 7/18 - (265/98)b_8\sqrt{21} - (215/14)b_8; \\
a_{11,8} &= (265/98)b_8\sqrt{21} + (215/14)b_8.
\end{aligned}$$

The numerical solution is given by the formula (1.2). The values of k_s are given by the formula (1.3). We can notice that if $b_8 = 49/180$ and $a_{10,5} = 1/9$, then we find the method of Cooper–Verner [1, 12].

With the help of Maple, the stability function depends on $a_{10,5}$ and is given by [12]:

$$\begin{aligned}
\zeta(z) &= 1 + z + \frac{1}{2}z^2 + \frac{1}{6}z^3 + \frac{1}{24}z^4 + \frac{1}{120}z^5 + \frac{1}{720}z^6 + \frac{1}{5040}z^7 + \frac{1}{40320}z^8 \\
&+ \left(-\frac{797}{50803200} + \frac{1}{25200}a_{10,5} + \frac{37}{4233600}\sqrt{21}a_{10,5} - \frac{499}{152409600}\sqrt{21} \right) z^9 \\
&+ \left(\frac{1}{470400} + \frac{1}{2083725}\sqrt{21} - \frac{31}{940800}a_{10,5} - \frac{61}{8467200}\sqrt{21}a_{10,5} \right) z^{10} \\
&+ \left(-\frac{1}{29030400} - \frac{13}{4267468800}\sqrt{21} + \frac{11}{1612800}a_{10,5} + \frac{353}{237081600}\sqrt{21}a_{10,5} \right) z^{11}.
\end{aligned}$$

For $a_{10,5} = 10^6$ we find

$$\begin{aligned} \zeta(z) = & 1 + z + \frac{1}{2}z^2 + \frac{1}{6}z^3 + \frac{1}{24}z^4 + \frac{1}{120}z^5 + \frac{1}{720}z^6 + \frac{1}{5040}z^7 \\ & + \frac{1}{40320}z^8 + \frac{2015999203}{50803200}z^9 - \frac{15499999}{470400}z^{10} + \frac{197999999}{29030400}z^{11} \\ & + \frac{190285643}{21772800}\sqrt{21}z^9 - \frac{60046871}{8334900}\sqrt{21}z^{10} + \frac{6353999987}{4267468800}\sqrt{21}z^{11}. \end{aligned}$$

The stability region of the new RK8 method for $a_{10,5} = 10^6$ is given by Fig. 3.




Figure 3. Stability region of the new RK8 method for $a_{10,5} = 10^6$.

We can see that the stability region of the new method of order 8 is smaller than 2, 3, 4. There is a decrease in the values of x and y .

For $a_{10,5} = 10^{12}$ the stability region is the following (see Fig. 4):




Figure 4. Stability region of the new RK8 method for $a_{10,5} = 10^{12}$.

We can see that the stability region of the new method of order 8 is smaller than those of ordering regions 1, 2, 3, 4. There is a decrease in the values of x and y .

For $a_{10,5} = \underbrace{9 \dots 9}_{37 \text{ times}}$ the stability region is shown at the next figure:




Figure 5. Stability region of the new RK8 method.

We find that the values of x and y have very strongly diminished and the region of stability is very small.

4. Conclusion

Presumably, by representing the domains of stability of methods of the order of 1, 2, 3, 4, one could assume that the higher the order, the greater the area of stability. However, a new 8 order method is discovered. The stability region of this 8 order method is smaller than that of the regions of orders 2, 3, 4. We can therefore conclude that the evolution of the size of the stability regions of Runge-Kutta methods does not depend on the order of the method.

REFERENCES

1. Butcher J.-C. *Numerical Methods for Ordinary Differential Equations*. 2nd ed. John Wiley & Sons Ltd., 2008. 175 p. DOI: [10.1002/9780470753767](https://doi.org/10.1002/9780470753767)
2. Calvo M., Montijano J. I., Randez L. A new embedded pair of Runge–Kutta formulas of orders 5 and 6. *Comput. Math. Appl.*, 1990. Vol. 20, No. 1. P. 15–24. DOI: [10.1016/0898-1221\(90\)90064-Q](https://doi.org/10.1016/0898-1221(90)90064-Q)
3. Cassity C. R. The complete solution of the fifth order Runge–Kutta equations. *SIAM J. Numer. Anal.*, 1969. Vol. 6, No. 3. P. 432–436. DOI: [10.1137/0706038](https://doi.org/10.1137/0706038)
4. Feagin T. A tenth-order Runge–Kutta method with error estimate. In: *Proc. of the IAENG Conf. on Scientific Computing*. Hong Kong, 2007. Accessible at <https://sce.uhcl.edu/feagin/courses/rk10.pdf>
5. Feagin T. *High-Order Explicit Runge-Kutta Methods*. 2013. Accessible at <http://sce.uhcl.edu/rungekutta>
6. Hairer E., Nørsett S. P., Wanner G. *Solving Ordinary Differential Equations I. Nonstiff Problems*. Springer Ser. Comput. Math., vol. 8. Berlin, Heidelberg: Springer-Verlag, 1993. 528 p. DOI: [10.1007/978-3-540-78862-1](https://doi.org/10.1007/978-3-540-78862-1)
7. Houben S. *Stability Regions of Runge–Kutta Methods*. Eindhoven University of Technology, 2002. Accessible at https://www.win.tue.nl/casa/meetings/seminar/previous/_abstract020220_files/talk.pdf
8. Jackiewicz Z. *General Linear Methods for Ordinary Differential Equations*. John Wiley & Sons, Inc., 2009. 482 p. DOI: [10.1002/9780470522165](https://doi.org/10.1002/9780470522165)
9. Khashin S. I. *List of Some Known Runge-Kutta Methods Family*. Preliminary version. 2013. Accessible at http://math.ivanovo.ac.ru/dalgebra/Khashin/rk/sh_rk.html
10. Kashin S. I. Estimating the error in classical Runge–Kutta methods. *Comput. Math. Math. Phys.*, 2014. Vol. 54, No. 5. P. 767–774. DOI: [10.1134/S0965542514050145](https://doi.org/10.1134/S0965542514050145)

-
11. Liu M.Z., Song M.H., Yang Z.W. Stability of Runge–Kutta methods in the numerical solution of equation $u'(t) = au(t) + a_0u([t])$. *J. Comput. Appl. Math.*, 2004. Vol. 166, No. 2. P. 361–370. DOI: [10.1016/j.cam.2003.04.002](https://doi.org/10.1016/j.cam.2003.04.002)
 12. Seka H., Assui K.R. A New Eighth Order Runge-Kutta Family Method. *J. Math. Res.*, 2019. Vol. 11, No. 2. P. 190–199. DOI: [10.5539/jmr.v11n2p190](https://doi.org/10.5539/jmr.v11n2p190)
 13. Velagala S.R. *Stability Analysis of the 4th order Runge–Kutta Method in Application to Colloidal Particle Interactions*. Master's thesis. University of Illinois, Urbana-Champaign, USA, 2014. Accessible at <http://hdl.handle.net/2142/72750>